

Invariance of the quasilinear equations of hiperbolic type with respect to the three-parametric Lie algebras

Olena Magda*

Institute of Mathematics of the Academy of Sciences of Ukraine,
Tereshchenkivska Street 3, 01601 Kyiv-4, Ukraine

Abstract

We have solved completely the problem of the description of quasi-linear hyperbolic differential equations in two independent variables that are invariant under three-parameter Lie groups.

The problem of group classification of differential equations is one of the central problems of modern symmetry analysis of differential equations[1]. One of the important classes of hyperbolic equations. The problem of group classification of such equations has been discussed by many authors (see for instance [2–9]). On this report we consider the problem of the group classification of equations of form:

$$u_{tt} = u_{xx} + F(t, x, u, u_x), \quad (1)$$

where $u = u(t, x)$ and F is an arbitrary nonlinear differentiable function, with $F_{u_x, u_x} \neq 0$ is an arbitrary nonlinear smooth function, which dependet variables u or u_x . We use following notation $u_x = \frac{\partial u}{\partial x}$, $u_{xx} = \frac{\partial^2 u}{\partial x^2}$, $F_{u_x} = \frac{\partial F}{\partial u_x}$, $u_t = \frac{\partial u}{\partial t}$, $u_{tt} = \frac{\partial^2 u}{\partial t^2}$. For the group classification of equation (1) we use the approach proposed in [10]. Here we give three fundamental results (for details, the reader is refered to [11]).

Theorem 1 *The infinitesimal operator of the symmetry group of the equation (1) has following form:*

$$X = (\lambda t + \lambda_1)\partial_t + (\lambda x + \lambda_2)\partial_x + (h(x)u + r(t, x))\partial_u, \quad (2)$$

where $\lambda, \lambda_1, \lambda_2$ are arbitrary real constants and $h(x), r(t, x)$ are arbitrary functions which satisfy the condition

$$\begin{aligned} r_{tt} - \frac{d^2 h}{dx^2} u - r_{xx} + (h - 2\lambda)F - (\lambda t + \lambda_1)F_t - (\lambda x + \lambda_2)F_x - \\ - (hu + r)F_u - 2u_x \frac{dh}{dx} - u_x(h - \lambda)F_{u_x} - \frac{dh}{dx} u F_{u_x} - r_x F_{u_x} = 0. \end{aligned} \quad (3)$$

*e-mail: magda@imath.kiev.ua

Theorem 2 *The equivalence group of the equation (1) is given by transformations of the following form:*

$$\bar{t} = \gamma t + \gamma_1, \quad \bar{x} = \epsilon \gamma x + \gamma_2, \quad v = \rho(x)u + \theta(t, x), \quad \gamma \neq 0, \quad \rho \neq 0, \quad \epsilon = \pm 1. \quad (4)$$

Using theorem 2, one can prove the following result:

Theorem 3 *In the class of operators (2), there are no realizations of the algebras $so(3)$ and $sl(2, R)$.*

From this theorem we obtain the following :

Note 1 *in the class of operators (2) there are no realizations of any real semi-simple Lie algebras;*

Note 2 *there are no equations (1) which has algebras of invariance, which are isomorphic by real semi-simple algebras, or conclude those algebras as subalgebras.*

The set of three-dimensional solvable Lie algebras consists of the following two decomposable Lie algebras:

$$\begin{aligned} A_{3.1} &= A_1 \oplus A_1 \oplus A_1 = 3A_1; \\ A_{3.2} &= A_{2.2} \oplus A_1, \quad [e_1, e_2] = e_2, \end{aligned}$$

and the following eight of non-decomposable Lie algebras:

$$\begin{aligned} A_{3.3} : \quad & [e_2, e_3] = e_1; \\ A_{3.4} : \quad & [e_1, e_3] = e_1, \quad [e_2, e_3] = e_1 + e_2; \\ A_{3.5} : \quad & [e_1, e_3] = e_1, \quad [e_2, e_3] = e_2; \\ A_{3.6} : \quad & [e_1, e_3] = e_1, \quad [e_2, e_3] = -e_2; \\ A_{3.7} : \quad & [e_1, e_3] = e_1, \quad [e_2, e_3] = qe_2, \quad (0 < |q| < 1); \end{aligned}$$

We give the realizations of the algebras $A_{3.3}, A_{3.4}, A_{3.5}, A_{3.6}, A_{3.7}, A_{3.8}, A_{3.9}$ and the corresponding values of the functions F in the equation (1)

$$\begin{aligned} A_{3.3}^1 &= \left\langle u\partial_u, \partial_t + \beta\partial_x, \frac{1}{\beta}xu\partial_u, \right\rangle, \beta > 0 : \\ &F = -u^{-1}u_x^2 + u\tilde{G}(\omega), \quad \omega = x - \beta t; \\ A_{3.3}^2 &= \langle u\partial_u, \partial_x, m\partial_t + xu\partial_u, \rangle, m \neq 0 : \\ &F = -u^{-1}u_x^2 + u\tilde{G}(\omega), \quad \omega = t - mu_xu^{-1}; \\ A_{3.3}^3 &= \langle \partial_u, \partial_x, m\partial_t + x\partial_u, \rangle, m \neq 0 : \\ &F = \tilde{G}(\omega), \quad \omega = mu_x - t; \\ A_{3.3}^4 &= \langle \partial_u, \partial_t, \partial_x + t\partial_u, \rangle : \\ &F = \tilde{G}(u_x); \\ A_{3.3}^5 &= \langle u\partial_u, \partial_x, xu\partial_u, \rangle : \end{aligned}$$

$$\begin{aligned}
& F = -u^{-1}u_x^2 + u\tilde{G}(t); \\
A_{3.3}^6 &= \langle t\partial_t + x\partial_x, u\partial_u, \ln|x|u\partial_u \rangle : \\
& F = -u^{-1}u_x^2 + \frac{1}{2}x^{-1}u_x + x^{-2}u\tilde{G}(\xi), \xi = tx^{-1}; \\
A_{3.3}^7 &= \left\langle u\partial_u, \partial_t + k\partial_x, m\partial_t + \frac{1}{k}xu\partial_u \right\rangle, k > 0, m \in R : \\
& F = -u^{-1}u_x^2 + u\tilde{G}(\omega), \omega = x - kt + mku^{-1}u_x; \\
A_{3.3}^8 &= \langle e^{mt}\partial_u, \partial_x, \partial_t + xe^{mt}\partial_u \rangle, m \neq 0 : \\
& F = m^2u + \tilde{G}(\omega), \omega = \frac{e^{mt}}{m} - u_x; \\
A_{3.3}^9 &= \langle \partial_u, \partial_t, t\partial_u \rangle : \\
& F = \tilde{G}(x, u_x); \\
A_{3.3}^{10} &= \langle u\partial_u, \partial_t - \beta^{-1}xu\partial_u, \partial_t + \beta\partial_x \rangle, \beta > 0 : \\
& F = -u^{-1}u_x^2 + u\tilde{G}(\omega), \omega = x - \beta t - \beta^2u_xu^{-1}; \\
A_{3.3}^{11} &= \langle u\partial_u, \partial_t - xu\partial_u, \partial_x \rangle : \\
& F = t^2u + 2tu_x + u\tilde{G}(\omega), \omega = t + u_xu^{-1}; \\
A_{3.3}^{12} &= \langle e^{kt}\partial_u, \partial_t + ku\partial_u, \beta\partial_x + te^{kt}\partial_u \rangle, \beta > 0, k > 0 : \\
& F = k^2u + \frac{2kx}{\beta}e^{kt} + e^{kt}\tilde{G}(\omega), \omega = e^{-kt}u_x; \\
A_{3.3}^{13} &= \langle |t|^{\frac{1}{2}}\partial_u, -|t|^{\frac{1}{2}}\ln|t|\partial_u, t\partial_t + x\partial_x + \frac{1}{2}u\partial_u \rangle : \\
& F = -\frac{u}{4}t^{-2} + u_x^3\tilde{G}(\omega, v), \omega = tx^{-1}, v = xu_x^2; \\
A_{3.3}^{14} &= \langle \partial_u, -t\partial_u, \partial_t + k\partial_x \rangle, k > 0 : \\
& F = \tilde{G}(\omega, u_x), \omega = x - kt; \\
A_{3.4}^1 &= \langle \partial_u, \partial_t + k\partial_x, t\partial_t + x\partial_x + (u+t)\partial_u \rangle, \eta = x - kt, m = 1, k > 0 : \\
& F = \eta^{-1}\tilde{G}(u_x); \\
A_{3.4}^2 &= \langle \eta^{m-1}\partial_u, \partial_t + k\partial_x, t\partial_t + x\partial_x + (mu + t\eta^{m-1})\partial_u \rangle, \\
& \eta = x - kt, k > 0, m \in R, m \neq 1, 2 : \\
& F = (k^2 - 1)(m - 1)(m - 2)\eta^{-2}u - \frac{2k(1 - m)}{2m - 4}\eta^{m-2} + \eta^{2-m}\tilde{G}(\omega), \\
& \omega = ((1 - m)u + \eta u_x)\eta^{3m-4}; \\
A_{3.4}^3 &= \langle \partial_u, \partial_t, t\partial_t + x\partial_x + (u+t)\partial_u \rangle : \\
& F = x^{-1}\tilde{G}(u_x); \\
A_{3.4}^4 &= \left\langle e^{ktx^{-1}}\partial_u, \partial_t + kx^{-1}u\partial_u, t\partial_t + x\partial_x + (u + te^{ktx^{-1}})\partial_u \right\rangle, k \neq 0 : \\
& F = u(k^2t^2x^{-4} - 2ktx^{-3} + k^2x^{-2}) + 2ktu_xx^{-2} + e^{ktx^{-1}}(2k\ln|x|x^{-1} + x^{-1}\tilde{G}(\omega)), \\
& \omega = e^{-ktx^{-1}}(u_x + ktux^{-2}); \\
A_{3.4}^5 &= \langle kx^{-1}u\partial_u, -k\ln|x|x^{-1}u\partial_u, t\partial_t + x\partial_x \rangle, k > 0 : \\
& F = -u^{-1}u_x^2 + 3xu_x + u\ln|u|x^{-2} + ux^{-2}\tilde{G}(\omega), \omega = tx^{-1};
\end{aligned}$$

$$\begin{aligned}
A_{3.4}^6 &= \langle kx^{\frac{1}{2}}u\partial_u, -k\ln|x|x^{\frac{1}{2}}u\partial_u, 2(t\partial_t + x\partial_x) \rangle, k > 0 : \\
&F = -u^{-1}u_x^2 + \frac{1}{4}u\ln|u|x^{-2} + ux^{-2}\tilde{G}(\omega), \omega = tx^{-1}; \\
A_{3.4}^7 &= \langle -\partial_t + 2\partial_x, e^{\frac{1}{2}x}u\partial_u, e^{\frac{1}{2}x}\partial_u \rangle : \\
&F = -u^{-1}u_x^2 - u_x + \frac{1}{4}u\ln|u| + u\tilde{G}(\eta), \eta = -2t - x; \\
A_{3.4}^8 &= \langle -2\partial_x, e^{\frac{1}{2}x}u\partial_u, e^{\frac{1}{2}x}xu\partial_u \rangle : \\
&F = -u^{-1}u_x^2 - u_x + \frac{1}{4}u\ln|u| + u\tilde{G}(t); \\
A_{3.4}^9 &= \langle kx^{-1}u\partial_u, \partial_t - k\ln|x|x^{-1}u\partial_u, t\partial_t + x\partial_x \rangle, k > 0 : \\
&F = k^2t^2ux^{-4} - 3ktux^{-3} + 2ktu_x x^{-2} + 2ktux^{-3}\ln|u| \\
&- 2ux^{-2}\ln|u| + 2u_x x^{-1}\ln|u| + x^{-2}u\ln^2|u| + ux^{-2}\tilde{G}(\omega), \\
&\omega = xu_xu^{-1} + \ln|u| + ktx^{-1} \\
A_{3.4}^{10} &= \langle |t|^{\frac{1}{2}}\partial_u, -|t|^{\frac{1}{2}}\ln|t|\partial_u, t\partial_t + x\partial_x + \frac{3}{2}u\partial_u \rangle : \\
&F = -\frac{u}{4}t^{-2} + u_x^{-1}\tilde{G}(\omega, v), \omega = tx^{-1}, v = x^{-1}u_x^2; \\
A_{3.4}^{11} &= \langle \partial_u, -t\partial_u, \partial_t + k\partial_x + u\partial_u \rangle, k > 0 : \\
&F = u_x\tilde{G}(\omega, v), \omega = x - kt, v = \ln|u_x| - t; \\
A_{3.5}^1 &= \langle \partial_u, \partial_t + k\partial_x, t\partial_t + x\partial_x + u\partial_u \rangle, \eta = x - kt, k > 0 : \\
&F = \zeta^{-1}\tilde{G}(u_x); \\
A_{3.5}^2 &= \langle \eta\partial_u, \partial_t + k\partial_x, t\partial_t + x\partial_x + 2u\partial_u \rangle, \eta = x - kt, k > 0 : \\
&F = \tilde{G}(\omega), \omega = (-u + u_x\eta)\eta^{-2}; \\
A_{3.5}^3 &= \langle \eta^{m-1}\partial_u, \partial_t + k\partial_x, t\partial_t + x\partial_x + mu\partial_u \rangle, \eta = x - kt, k > 0, m \in R, m \neq 1, 2 : \\
&F = (k^2 - 1)(m - 1)(m - 2)u\eta^{-2} + \eta^{m-2}\tilde{G}(\omega), \omega = ((1 - m)u + u_x\eta)\eta^{-m}; \\
A_{3.5}^4 &= \langle \partial_x, \partial_u, t\partial_t + x\partial_x + u\partial_u \rangle : \\
&F = t^{-1}\tilde{G}(u_x); \\
A_{3.5}^5 &= \langle \partial_x, t\partial_u, t\partial_t + x\partial_x + 2u\partial_u \rangle : \\
&F = \tilde{G}(\omega), \omega = u_x t; \\
A_{3.5}^6 &= \langle \partial_x, |t|^{m-1}\partial_u, t\partial_t + x\partial_x + mu\partial_u \rangle, m \neq 1, 2, m \in R : \\
&F = (2u - 3mu - m^2u)t^{-2} + t^{m-2}\tilde{G}(\omega), \omega = u_x t^{m-1}; \\
A_{3.5}^7 &= \langle \partial_t, \partial_x, t\partial_t + x\partial_x \rangle : \\
&F = u_x^2\tilde{G}(u); \\
A_{3.5}^8 &= \langle \partial_t, \partial_x, t\partial_t + x\partial_x + mu\partial_u \rangle, m \neq 0, 1, 2 : \\
&F = |u|^{\frac{m-2}{m}}\tilde{G}(\omega), \omega = u_x^{-1}|u|^{\frac{m-1}{m}}; \\
A_{3.5}^9 &= \langle \partial_t, \partial_x, t\partial_t + x\partial_x + \partial_u \rangle, m \neq 0 : \\
&F = e^{-2u}\tilde{G}(\omega), \omega = e^u u_x; \\
A_{3.5}^{10} &= \langle \partial_t, x^{-1}u\partial_u, t\partial_t + x\partial_x \rangle, k \neq 0 :
\end{aligned}$$

$$\begin{aligned}
& F = 2u_x x^{-1} \ln |u| + u \ln^2 |u| x^{-2} - 2u \ln |u| x^{-2} + x^{-2} u \tilde{G}(\omega), \\
& \omega = u_x u^{-1} x + \ln |u|; \\
A_{3.5}^{11} &= \langle \partial_t + kx^{-1} u \partial_u, e^{ktx^{-1}} \partial_u, t\partial_t + x\partial_x + u\partial_u \rangle, \quad k \neq 0 : \\
& F = uk(kt^2 - 2xt + kx^2)x^{-4} + 2ktu_x x^{-2} + e^{ktx^{-1}} x^{-1} \tilde{G}(\omega), \\
& \omega = e^{-ktx^{-1}} (u_x + ktu x^{-2}); \\
A_{3.5}^{12} &= \langle \partial_t, \partial_u, t\partial_t + x\partial_x + u\partial_u \rangle : \\
& F = x^{-1} \tilde{G}(u_x); \\
A_{3.6}^1 &= \langle \partial_t + k\partial_x, \partial_u, t\partial_t + x\partial_x - u\partial_u \rangle, \eta = x - kt, \quad k > 0 : \\
& F = \eta^3 \tilde{G}(\omega), \quad \omega = \eta^{-2} u_x; \\
A_{3.6}^2 &= \langle \partial_t + k\partial_x, \eta \partial_u, t\partial_t + x\partial_x \rangle, \eta = x - kt, \quad k > 0 : \\
& F = \eta^2 \tilde{G}(\omega), \quad \omega = -u + \eta u_x; \\
A_{3.6}^3 &= \langle \partial_t + k\partial_x, \eta^{m+1} \partial_u, t\partial_t + x\partial_x + mu\partial_u \rangle, \eta = x - kt, \quad k > 0, \quad m \in R, \quad m \neq 0, -1 : \\
& F = (k^2 - 1)m(m+1)u\eta^{-2} + \eta^{2-m} \tilde{G}(\omega), \quad \omega = (-(m+1)u + \eta u_x) \eta^{3m}; \\
A_{3.6}^4 &= \langle \partial_u, \partial_x, -t\partial_t - x\partial_x + u\partial_u \rangle : \quad F = t^{-3} \tilde{G}(\omega), \quad \omega = u_x t^2; \\
A_{3.6}^5 &= \langle t\partial_u, \partial_x, -t\partial_t - x\partial_x + 2u\partial_u \rangle : \quad F = t^{-4} \tilde{G}(\omega), \quad \omega = u_x t^3; \\
A_{3.6}^6 &= \langle t^{m-1} \partial_u, \partial_x, -t\partial_t - x\partial_x + mu\partial_u \rangle, \quad m \neq 1, 2, \quad m \in R : \\
& F = (2u - 3mu + m^2 u) t^{-2} + t^{-(m+2)} \tilde{G}(\omega), \quad \omega = u_x t^{m+1}; \\
A_{3.6}^7 &= \langle \partial_x, t^{m+1} \partial_u, t\partial_t + x\partial_x + mu\partial_u \rangle, \quad m \neq 0, -1, \quad m \in R : \\
& F = m(m+1)u t^{-2} + t^{m-2} \tilde{G}(\omega), \quad \omega = u_x t^{1-m}; \\
A_{3.6}^8 &= \langle \partial_x, t\partial_u, t\partial_t + x\partial_x \rangle : \\
& F = t^{-2} \tilde{G}(\omega), \quad \omega = u_x t; \\
A_{3.6}^9 &= \langle \partial_x, \partial_u, t\partial_t + x\partial_x - u\partial_u \rangle : \\
& F = t^{-3} \tilde{G}(\omega), \quad \omega = u_x t^2; \\
A_{3.6}^{10} &= \langle \partial_t, \partial_u, t\partial_t + x\partial_x - u\partial_u \rangle : \\
& F = x^{-2} \tilde{G}(\omega), \quad \omega = u_x x^2; \\
A_{3.6}^{11} &= \langle \partial_t, xu\partial_u, t\partial_t + x\partial_x \rangle, \quad k \neq 0 : \\
& F = -2u_x x^{-1} \ln |u| + u \ln^2 |u| x^{-2} + x^{-2} u \tilde{G}(\omega), \\
& \omega = u_x u^{-1} x - \ln |u|; \\
A_{3.6}^{12} &= \langle \partial_t + mx^{-1} u \partial_u, xu\partial_u, t\partial_t + x\partial_x \rangle, \quad mk \neq 0 : \\
& F = 4m^2 t^2 x^{-4} u - 2mtu x^{-3} + 4mtu_x x^{-2} - 4mtu x^{-3} \ln |u| - \\
& 2u_x x^{-1} \ln |u| + u \ln^2 |u| x^{-2} + x^{-2} u \tilde{G}(\omega), \\
& \omega = u_x u^{-1} x - \ln |u| + 2mtx^{-1}; \\
A_{3.6}^{13} &= \langle \partial_t + kx^{-1} u \partial_u, e^{ktx^{-1}} \partial_u, t\partial_t + x\partial_x u \partial_u \rangle, \quad k \neq 0 : \\
& F = uk(kt^2 - 2xt + kx^2)x^{-4} + 2ktu_x x^{-2} + e^{ktx^{-1}} x^{-3} \tilde{G}(\omega), \\
& \omega = e^{-ktx^{-1}} (u_x x^2 + ktu); \\
A_{3.6}^{14} &= \langle kx^{-1} u \partial_u, mxu\partial_u, t\partial_t + x\partial_x \rangle, \quad mk \neq 0 :
\end{aligned}$$

$$\begin{aligned}
& F = -u^{-1}u_x^2 + xu_x - u \ln |u|x^{-2} + ux^{-2}\tilde{G}(\omega), \quad \omega = tx^{-1}; \\
A_{3.6}^{15} &= \langle -\partial_t + \partial_x, e^x u \partial_u, e^{-x} u \partial_u \rangle : \\
& F = -u^{-1}u_x^2 - u \ln |u| + u\tilde{G}(\eta), \quad \eta = -t - x; \\
A_{3.6}^{16} &= \langle \partial_x, e^x u \partial_u, e^{-x} u \partial_u \rangle : \\
& F = -u^{-1}u_x^2 - \ln |u| + u\tilde{G}(t); \\
A_{3.6}^{17} &= \langle e^{(m-1)t} \partial_u, e^{(1-m)t} \partial_u, \partial_t + k\partial_x + mu\partial_u \rangle, k \geq 0, m \neq 1 : \\
& F = u - 2mu - m^2u + u_x\tilde{G}(\omega, v), \quad \omega = x - kt, \quad v = \ln |u_x| - mt; \\
\\
A_{3.7}^1 &= \langle \partial_t + k\partial_x, \partial_u, t\partial_t + x\partial_x + qu\partial_u \rangle, \eta = x - kt, \quad k > 0, \quad 0 < |q| < 1 : \\
& F = \eta^{q-2}\tilde{G}(\omega), \quad \omega = \eta^{q-1}u_x^{-1}; \\
A_{3.7}^2 &= \langle \partial_t + k\partial_x, \eta\partial_u, t\partial_t + x\partial_x + (q+1)u\partial_u \rangle, \eta = x - kt, \quad k > 0, \quad 0 < |q| < 1 : \\
& F = \eta^{q-1}\tilde{G}(\omega), \quad \omega = \eta^{-q}(u_x - u\eta^{-1}); \\
A_{3.7}^3 &= \langle \partial_t + k\partial_x, \eta^{m-q}\partial_u, t\partial_t + x\partial_x + mu\partial_u \rangle, \\
& \eta = x - kt, \quad k > 0, \quad m \in R, m \neq q, q+1, \quad 0 < |q| < 1 : \\
& F = (k^2 - 1)(m - q)(m - q - 1)\eta^{-2}u + \eta^{m-2}\tilde{G}(\omega), \quad \omega = \eta^{1-m}(u_x - (m - q)u\eta^{-1}); \\
A_{3.7}^4 &= \langle \partial_x, \partial_u, t\partial_t + x\partial_x + qu\partial_u \rangle, \quad 0 < |q| < 1 : \quad F = t^{q-2}\tilde{G}(\omega), \quad \omega = u_x^{-1}t^{q-1}; \\
A_{3.7}^5 &= \langle \partial_x, t\partial_u, t\partial_t + x\partial_x + (q+1)u\partial_u \rangle, \quad 0 < |q| < 1 : \\
& F = t^{q-1}\tilde{G}(\omega), \quad \omega = u_x^{-1}t^q; \\
A_{3.7}^6 &= \langle \partial_x, t^{m-q}\partial_u, t\partial_t + x\partial_x + mu\partial_u \rangle, \quad m \neq q, q+1, \quad m \in R, \quad 0 < |q| < 1 : \\
& F = (m - 1 - q)(m - q)t^{-2}u + t^{m-2}\tilde{G}(\omega), \quad \omega = u_x^{-1}t^{m-1}; \\
A_{3.7}^7 &= \langle \partial_t, \partial_u, t\partial_t + x\partial_x + qu\partial_u \rangle, \quad 0 < |q| < 1 : \\
& F = x^{q-2}\tilde{G}(\omega), \quad \omega = u_x^{-1}x^{q-1}; \\
A_{3.7}^8 &= \langle \partial_t, x^{-q}u\partial_u, t\partial_t + x\partial_x \rangle, k \neq 0, \quad 0 < |q| < 1 : \\
& F = 2qu_x x^{-1} \ln |u| + q^2u \ln^2 |u|x^{-2} - (q+1)qx^{-2}u \ln |u| + x^{-2}u\tilde{G}(\omega), \\
& \omega = u_x u^{-1}x + q \ln |u|; \\
A_{3.7}^9 &= \langle \partial_t + mx^{-1}u\partial_u, kx^{-q}u\partial_u, t\partial_t + x\partial_x \rangle, mk \neq 0, \quad 0 < |q| < 1 : \\
& F = (m(q-1)t(m(q-1)tu + x((q+2)u - 2u_x x))x^{-4} - \\
& qx(2m(q-1)tu + x(u + qu - 2u_x x)) \ln |u|x^{-3} + q^2ux^{-2} \ln^2 |u| + ux^{-2}\tilde{G}(\omega), \\
& \omega = u_x ux + q \ln |u| + m(1-q)tx^{-1} \\
A_{3.7}^{10} &= \langle \partial_t + kx^{-1}u\partial_u, e^{ktx^{-1}}\partial_u, t\partial_t + x\partial_x + qu\partial_u \rangle, \quad k \neq 0, \quad 0 < |q| < 1 : \\
& F = uk(kt^2x^{-4} - 2tx + kx^{-2}) + 2ktu_x x^{-2} + e^{ktx^{-1}} + x^{q-2}\tilde{G}(\omega), \\
& \omega = e^{-ktx^{-1}}x^{1-q}(u_x + ktux^{-2}); \\
A_{3.7}^{11} &= \langle kx^{-1}u\partial_u, mx^{-q}u\partial_u, t\partial_t + x\partial_x \rangle, \quad mk \neq 0, \quad 0 < |q| < 1 : \\
& F = -u^{-1}u_x^2 + (q+2)x^{-1}u_x + qu \ln |u|x^{-1} + ux^{-2}\tilde{G}(\omega), \quad \omega = tx^{-1}; \\
A_{3.7}^{12} &= \langle -\partial_t + 2\partial_x, e^{\frac{1}{2}qx}u\partial_u, e^{\frac{1}{2}x}xu\partial_u \rangle, \quad 0 < |q| < 1 :
\end{aligned}$$

$$\begin{aligned}
& F = -u^{-1}u_x^2 - \frac{q+1}{2}u_x + \frac{q}{4}u \ln |u| + u\tilde{G}(\eta), \quad \eta = -2t - x; \\
A_{3.7}^{13} &= \langle -2\partial_x, e^{\frac{1}{2}x}u\partial_u, e^{\frac{1}{2}qx}xu\partial_u \rangle, \quad 0 < |q| < 1 : \\
& F = -u^{-1}u_x^2 - \frac{q+1}{2}u_x + \frac{q}{4}u \ln |u| + u\tilde{G}(t); \\
A_{3.7}^{14} &= \langle e^{\frac{1}{2}(q+1)t}\partial_u, e^{\frac{1}{2}(1-q)t}\partial_u, \partial_t + k\partial_x + \frac{1}{2}(1+q)u\partial_u \rangle, \quad k \geq 0, \quad 0 < |q| < 1 : \\
& F = \frac{(q+1)^2}{4}u + u_x\tilde{G}(\omega, v), \quad \omega = x - kt, \quad v = \ln |u_x| - \frac{q+1}{2}t; \\
A_{3.8}^1 &= \langle \cos(\ln |x|)u\partial_u, \sin(\ln |x|)u\partial_u, t\partial_t + x\partial_x \rangle : \\
& F = -u^{-1}u_x^2 + u_x x^{-1} + u \ln |u|x^{-2} + ux^{-2}\tilde{G}(\omega), \quad \omega = tx^{-1}; \\
A_{3.8}^2 &= \langle \cos xu\partial_u, \sin xu\partial_u, \partial_x \rangle : \\
& F = -u^{-1}u_x^2 + u \ln |u| + u\tilde{G}(t); \\
A_{3.8}^3 &= \langle \cos xu\partial_u, \sin xu\partial_u, \partial_t + \partial_x \rangle : \\
& F = -u^{-1}u_x^2 + u \ln |u| + u\tilde{G}(\eta), \quad \eta = x - t; \\
A_{3.8}^4 &= \langle \cos t\partial_u, -\sin t\partial_u, \partial_t + k\partial_x \rangle, \quad k \geq 0 : \\
& F = -u + \tilde{G}(\eta, u_x), \quad \eta = x - kt; \\
A_{3.8}^5 &= \left\langle |t|^{\frac{1}{2}} \cos\left(\frac{\ln |t|}{2k}\right) \partial_u, -|t|^{\frac{1}{2}} \sin\left(\frac{\ln |t|}{2k}\right) \partial_u, k(2t\partial_t + 2x\partial_x + u\partial_u) \right\rangle, \quad k \geq 0 : \\
& F = -\frac{k^2+1}{4k^2}t^{-2}u + |t|^{-\frac{1}{2}}\tilde{G}(\omega, v), \quad \omega = tx^{-1}, \quad v = tu_x^2; \\
A_{3.9}^1 &= \langle x^{(q^2+1)} \cos((q^2+1) \ln |x|)u\partial_u, x^{(q^2+1)} \sin((q^2+1) \ln |x|)u\partial_u, t\partial_t + x\partial_x \rangle, \quad q > 0 : \\
& F = -u^{-1}u_x^2 + (2q^2+1)u_x x^{-1} + 2(q^2+1)^2u \ln |u|x^{-2} + ux^{-2}\tilde{G}(\omega), \quad \omega = tx^{-1}; \\
A_{3.9}^2 &= \langle e^{\frac{1}{2}qx} \cos \frac{x}{2}u\partial_u, e^{\frac{1}{2}qx} \sin \frac{x}{2}u\partial_u, -2\partial_x \rangle, \quad q > 0 : \\
& F = -u^{-1}u_x^2 - qu_x + \frac{(q^2+1)u \ln |u|}{4} + u\tilde{G}(t); \\
A_{3.9}^3 &= \langle e^{\frac{1}{2}qx} \cos \frac{x}{2}u\partial_u, e^{\frac{1}{2}qx} \sin \frac{x}{2}u\partial_u, -\partial_t + 2\partial_x \rangle, \quad q > 0 : \\
& F = -u^{-1}u_x^2 - qu_x + \frac{(q^2+1)u \ln |u|}{4} + u\tilde{G}(\eta), \quad \eta = -2t - x; \\
A_{3.9}^4 &= \langle \sin t\partial_u, \cos t\partial_u, \partial_t + k\partial_x + qu\partial_u \rangle, \quad k \geq 0, \quad q > 0 : \\
& F = -u + u_x\tilde{G}(\eta, v), \quad \eta = x - kt, \quad v = e^{-qt}u_x; \\
A_{3.9}^5 &= \left\langle |t|^{\frac{1}{2}} \sin\left(\frac{\ln |t|}{2(k-q)}\right) \partial_u, |t|^{\frac{1}{2}} \cos\left(\frac{\ln |t|}{2(k-q)}\right) \partial_u, 2(k-q)(t\partial_t + x\partial_x) + ku\partial_u \right\rangle, \\
& k \in R, \quad q > 0, \quad k \neq q : \\
& F = -\frac{(k-q)^2+1}{4(k-q)^2}t^{-2}u + |t|^{\frac{4q-3k}{2(k-q)}}\tilde{G}(\omega, v), \quad \omega = tx^{-1}, \quad v = |t|^{k-2q}|u_x|^{2(k-q)};
\end{aligned}$$

Acknowledgements

The author is grateful to R.Z. Zhdanov and V.I. Lahno for proposing of the problem and for the help in the research.

References

- [1] Ovsjannikov L.V. Group Analysis of Differential Equations, Nauka, Moscow, 1978
- [2] Ames W.F., Lochner R.I., and Adams E. Int. Nonlin. Mech.16, 439 (1981)
- [3] Torrisi M. and Valenti A. Int. J. Nonlin. Mech. 20, 139 (1985)
- [4] Torrisi M. and Valenti A. Atteisem. Math. Fis. Uniu. Modena XXXVIII, 445 (1990)
- [5] Ibragimov N.H. Torrisi M. and Valenti A. J. Math. Phys. 32, 2988 (1991)
- [6] Oron A. and Rosenau Ph. Phys. Lett. A. 118, 172 (1986)
- [7] Pucci E. and Salvatori M.C. Group properties of a class of semilinear hyperbolic equations // Int. J. Non. Mech. — 1986. — **21**, 2. — P. 147–152.
- [8] Pucci E. Group analysis of the equation $u_{tt} + \lambda u_{xx} = g(u, u_x)$ // Riv. Mat. Univ. Parma. — 1987. — **12**, 4. — P. 71–87.
- [9] Ames W.F., Adams E., Lohner R.J. Group properties of $u_{tt} = [f(u)u_x]_x$ // Int. J. Non. Mech. — 1981. — **16**, 5–6. — P. 439–447.
- [10] Zhdanov R.Z., Lahno V.I. Group classification of heat conductivity equations with a nonlinear source // J. Phys. A: Math. Gen. — 1999. — **32**. — 7405–7418.
- [11] Lahno V., Magda O., Zhdanov R. On invariance quasi-linear hyperbolic-type differential equations under three-dimensional Lie algebras // Proceedings of Institute of Mathematics of NAS of Ukraine, T.36 / Eds.: A.G. Nikitin, 2001. 316p